Table 6: Monitoring Indicators

Table 6: Comparable maternal-fetal indicators available on AHS tableau

| | MANION CONTINUE TO STATE OF THE | |
|---|--|--|
| | MNCY SCN monitoring indicator | Comparable existing indicator in AHS tableau |
| 1 | Caesarean section rate without medical indication (e.g., labour, others) by gestational age | Obstetric delivery and newborn dashboard (birth site based): cesarean section, vaginal birth after c-section (VBAC), and induction If interested in this indicator, email seija.kromm@ahs.ca. |
| 2 | Proportion of women with prior caesarean section attempting vaginal birth with a trial of labour (successful/ unsuccessful) | Obstetric delivery and newborn dashboard (birth site based): cesarean section, vaginal birth after c-section (VBAC), and induction If interested in this indicator, email seija.kromm@ahs.ca . |
| 3 | Rate of induction of labour without medical indication by gestational age (GA) | Obstetric delivery and newborn dashboard (birth site based): cesarean section, vaginal birth after c-section (VBAC), and induction If interested in this indicator, email seija.kromm@ahs.ca. Obstetric delivery and newborn dashboard (birth site based): cesarean section, vaginal birth after c-section (VBAC), and induction by GA If interested in this indicator, email seija.kromm@ahs.ca. |
| 4 | Screening rate for postpartum depression (PPD) following delivery within first year postpartum | Percentage of mothers Screened for PPD (EPDS Score 13+) If interested in this indicator, email seija.kromm@ahs.ca. |
| 5 | Proportion of women with a positive screen for PPD (greater than or equal to 13 on the Edinburgh postnatal depression scale) within first year postpartum and referred | Positive screen for PPD (>=13) and referred If interested in this indicator, email <u>seija.kromm@ahs.ca</u>. |

February 2020



