

Measles, Mumps, Rubella, Varicella (MMR-Var) Vaccine

Get protected, get immunized.

- Vaccines make your immune system stronger. They build antibodies to help prevent diseases.
- Immunization is safe. It is much safer to get immunized than to get these diseases.

What is the MMR-Var vaccine?

The MMR-Var vaccine protects against measles, mumps, rubella, and varicella (chickenpox).

What is measles?

Measles is a virus that spreads easily through the air when someone who has measles coughs or sneezes. It can cause:

- a high fever
- a cough
- a runny nose
- red eyes
- a blotchy, red rash

Measles can be dangerous because:

- 1 in 10 people with measles will get middle ear or lung infections
- 1 in 1,000 people will get encephalitis (swelling of the brain), which can lead to seizures, deafness, or brain damage
- 2 to 3 of every 1,000 people with measles could die

What is mumps?

Mumps is a virus that spreads by coughing, sneezing, or having contact with saliva (such as kissing or sharing toys). You can have no symptoms but still spread mumps. It can cause:

- a fever
- a headache
- swelling of the glands around your jaw
- swelling of the testicles or ovaries
- deafness
- encephalitis
- meningitis (infection of the fluid and lining that cover the brain and spinal cord)

What is rubella?

Rubella is a virus that spreads by coughing or sneezing. It is usually mild. It can cause:

- a fever
- a sore throat
- swollen neck glands
- a rash with red, raised bumps
- painful, swollen joints
- encephalitis
- a bleeding disorder

If you get rubella while you are pregnant, it can cause loss of a baby during pregnancy (miscarriage or stillbirth) or the baby may be born with disabilities.

What is varicella (chickenpox)?

Varicella is a virus that can cause a fever and an itchy rash that looks like small water-filled blisters. It spreads easily through the air by coughing, sneezing, or by touching open blisters.

It is usually mild, but can be more serious in newborns, adults, and people with weak immune systems. People who get varicella can have:

- skin infections
- pneumonia (a type of lung infection)
- blood infections
- other deadly infections

Who should get the MMR-Var vaccine?

As part of the routine immunization schedule, children get this vaccine starting at age 12 months up to and including 12 years. You do not get this vaccine if you are older than 12 years.

How many doses do I need?

Children need 2 doses. As of January 1, 2021, children can get their first dose at age 12 months and their second dose at age 18 months. If your child did not get an MMR-Var vaccine at age 18 months, they can get it at age 4 years.

Are there other vaccines that protect against measles, mumps, rubella, and varicella?

The MMR vaccine protects against measles, mumps, and rubella. The VZ vaccine protects against varicella.

How well does the vaccine work?

After 2 doses, protection is about:

- 100% for measles
- 76% to 95% for mumps
- 95% for rubella
- 98% for varicella

Where can I get the MMR-Var vaccine?

You can get the vaccine at your local public health or community health centre.

Are there side effects from the MMR-Var vaccine?

There can be side effects from the MMR-Var vaccine. They tend to be mild and go away in a few days, but side effects can happen up to 6 weeks after getting this vaccine. They may include:

- redness, swelling, bruising, rash, or feeling sore where you had the needle
- getting upset easily
- a fever
- vomiting (throwing up) or loose stool (diarrhea)
- a measles-like rash (a blotchy, red rash), rubella-like rash (rash with red, raised bumps), or varicella-like rash (rash with water-filled blisters)

At least 1 out of 100 people who got this vaccine reported 1 or more of these side effects. In some cases, it is unknown if the vaccine caused these side effects.

It is important to stay at the clinic for 15 minutes after your vaccine. Some people may have a rare but serious allergic reaction called anaphylaxis. If anaphylaxis happens, you will get medicine to treat the symptoms.

It is rare to have a serious side effect after a vaccine. Call Health Link at 811 to report any serious or unusual side effects.

How can I manage side effects?

- To help with soreness and swelling, put a cool, wet cloth over the area where your child had the needle.
- There is medicine to help with a fever or pain. Check with your child's doctor or pharmacist if you are not sure what medicine or dose to give. Follow the directions on the package.
- Children under the age of 18 years should **not** take aspirin for 6 weeks after getting a vaccine that protects against varicella (unless your doctor has told you to take it) because it can cause serious health problems.
- If your child gets a rash that looks like varicella, keep it covered. If you cannot cover your child's rash, keep them away from anyone who is pregnant, newborn babies, and people with weak immune systems. If your child has more than 50 spots, call Health Link at 811.
- Some people with health problems, such as a weak immune system, must call their doctor if they get a fever. If you have been told to do this, call your child's doctor even if you think the fever is from the vaccine.

Who should not get the MMR-Var vaccine?

Your child may not be able to get this vaccine if:

- They have an allergy to any part of the vaccine.
- They had a severe (serious) or unusual side effect after this vaccine or one like it.
- They have a weak immune system (because of medicine they take or a health problem).
- They have a family history of a weak immune system.
- They had a blood product in the past 11 months.
- They had another live vaccine in the past 3 months.
- They have seizures or a family history of seizures.
- They had an organ or stem cell transplant.
- They are pregnant.

If your child has allergies or has had a side effect from this vaccine in the past, check with your child's doctor or a public health nurse before they get the vaccine.

Although your child can get the vaccine if they have a mild illness such as a cold or fever, they should stay home until they are feeling better to prevent spreading their illness to others.

My child is afraid of needles. How can I help my child prepare for their immunization?

Many children are afraid of needles. You can do many things before, during, and after your child's immunization to help them be more comfortable.



Scan the QR code, visit ahs.ca/vaccinecomfort, or call Health Link at 811 for information.

For more information about immunization



Call Health Link at 811



Scan the QR code or go to ahs.ca/immunize